

Supplemental Material

**Src-Mediated EGF Receptor Activation Regulates Ozone-Induced
Interleukin 8 Expression in Human Bronchial Epithelial Cells**

Weidong Wu, Phillip A. Wages, Robert B. Devlin, David Diaz-Sanchez, David B. Peden, and
James M. Samet

Figure S1




Figure S1. Src kinase is required for O₃-induced EGFR phosphorylation. BEAS-2B cells grown to confluence were pretreated with vehicle control (0.1% DMSO), or the Src kinase inhibitor Bosutinib (5 or 10 μM) for 30 min prior to exposure to 1 ppm O₃ for 1h. Proteins were extracted from the cells and subjected to SDS-PAGE followed by immunoblotting using phospho-specific EGFR antibodies, followed by a pan-EGFR antibody. % P indicates the optical density of the p-EGFR band as a fraction of the total EGFR signal (p-EGFR+EGFR). Data shown are representative of three separate experiments. *, indicates $P < 0.05$ compared to O₃ in DMSO group.

Figure S2




Figure S2. Src kinase is required for Zn²⁺-induced phosphorylation of EGFR in BEAS-2B cells. BEAS-2B cells grown to confluence were pretreated with vehicle control (0.1% DMSO), or the Src kinase inhibitor Bosutinib (10 μ M) for 30 min prior to exposure to exposed to 100 μ M ZnSO₄ for 1 h. Extracted cellular proteins were subjected to SDS-PAGE followed by immunoblotting using phospho-specific EGFR antibodies, followed by a pan--EGFR antibody. % P indicates the optical density of the p-EGFR band as a fraction of the total EGFR signal (p-EGFR+EGFR). Data shown are representative of three separate experiments.